

**UNCLASSIFIED**

**AD - 4 2 6 4 8 5**

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

426485

CATALOGED BY DDC

AS AD No.

A GRADIENT INEQUALITY FOR  
NON-DIFFERENTIABLE FUNCTIONS



by  
E. Eisenberg

ORC 63-25 (RR)  
SEPTEMBER 1963

OPERATIONS RESEARCH CENTER

INSTITUTE OF ENGINEERING RESEARCH

UNIVERSITY OF CALIFORNIA - BERKELEY



A GRADIENT INEQUALITY FOR NON-DIFFERENTIABLE FUNCTIONS

by

E. Eisenberg  
Operations Research Center  
University of California, Berkeley

September 1963

ORC 63-25 (RR)

This research has been partially supported by the Office of Naval Research under Contract Nonr-222(83) with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

# A GRADIENT INEQUALITY FOR NON-DIFFERENTIABLE FUNCTIONS

## I. Introduction

In mathematical programming the following result is very useful:

### THEOREM 1

**Assumption:**  $K$  is a nonempty open subset of  $R^n$ ,  $f: K \rightarrow R$ ,  $f$  is differentiable on  $K$ ,  $X$  is a convex subset of  $K$ .

**Conclusion:**  $f$  is convex on  $X$  if, and only if,  $f(x) - f(y) \geq (x - y)^T f'(y)$  for all  $x, y \in X$ ,

where  $f'(y)$  denotes the gradient of  $f$  at  $y$ .

The above theorem is stated, in a slightly weaker form, in [8 . Vol. 1, p. 405]; for the sake of completeness we present a proof of Theorem 2 in Appendix A. A related result, extremely useful in programming theory, and discussed in [3] is:

### THEOREM 2

**Assumption:**  $K$  is an open subset of  $R^n$ ,  $f: K \rightarrow R$ ,  $f$  is differentiable on  $K$ ,  $X$  is a convex subset of  $K$ ,  $f$  is convex on  $X$ ,  $x_0 \in X$ .

**Conclusion:**  $f(x_0) \leq f(x)$  for all  $x \in X$  if, and only if,  $(x - x_0)^T f'(x_0) \geq 0$  for all  $x \in X$ .

---

$R^n$  denotes the set of all column  $n$ -tuples with real number components, we write  $R$  in place of  $R^1$ .

$\dagger\dagger$  At each point of  $K$ , all first partial derivatives exist.

The proof of Theorem 2 is straightforward; if we know that  $(x - x_0)^T f'(x_0) \geq 0$  whenever  $x \in X$  then a direct application of Theorem 1, with  $y = x_0$ , yields:  $f(x_0) \leq f(x)$  for all  $x \in X$ . Conversely, if  $x_0$  minimizes  $f$  on  $X$  then for each  $x \in X$  and  $\lambda \in (0, 1)$ , since  $\lambda x + (1 - \lambda)x_0 \in X$ , we must have:  $f(x_0) \leq f(\lambda x + (1 - \lambda)x_0) = f(x_0 + \lambda(x - x_0))$ . As a consequence we have:

$$\frac{f(x_0 + \lambda(x - x_0)) - f(x_0)}{\lambda} \geq 0 \quad \text{whenever } \lambda \in (0, 1),$$

letting  $\lambda$  approach zero we obtain  $(x - x_0)^T f'(x_0) \geq 0$ . Q.E.D.

We shall be concerned here with generalizing Theorem 2 by relaxing the hypothesis that  $f$  is differentiable everywhere on  $K$  (and thus we shall actually be able to omit mentioning  $K$ ), though keeping the requirement that  $f$  be convex on  $X$ . In case we know only that  $f$  is convex, we still wish to obtain nontrivial characterizations of the fact that  $f$  has a minimum on  $X$  and, if possible, obtain some information about the minimizing point (i.e.,  $x_0$ ). We will be able to obtain such characterizations when  $X$  and  $f$  are of a special form, though  $f$  may fail to be differentiable.

In what follows we shall frequently apply the so-called Minkowski-Farkas lemma and we list it here for reference:

Lemma 1: Let  $A$  be any real  $m \times n$  matrix and let  $b \in \mathbb{R}^m$ . The following statements are equivalent:

- (i)  $b^T \pi \leq 0$  whenever  $A^T \pi \leq 0$ ,  $\pi \in \mathbb{R}^m$ ,
- (ii) there exists an  $x \in \mathbb{R}^n$  such that  $x \geq 0$ ,  $Ax = b$

---

<sup>†</sup>A vector inequality means that the inequality indicated holds for each component.

## II. Discussion of Special Problem

Suppose we are given an  $n$ -tuple  $a = (a_1, \dots, a_n)$ , an  $n \times n$  real symmetric positive semi-definite matrix  $C$  and a real  $m \times n$  matrix  $A$ .

We define:

$$(1) \quad \begin{aligned} X &= R^n \cap \{x \mid Ax \leq 0\} \\ f(x) &= ax + (x^T C x)^{\frac{1}{2}}, \quad \text{for } x \in R^n \end{aligned}$$

It is immediate that  $X$  is a convex cone (it is in fact a "finite cone"), also  $X$  is obviously nonempty because  $0 \in X$ . In addition, it follows readily from a well known inequality [4, Theorem 1 - (ii)] that  $f$  is a convex function on  $R^n$ .

With  $f$  and  $x$  as defined in (1) and  $x_0 \in X$  we should like to obtain a statement similar to the conclusion in Theorem 2. Some direct considerations lead us to believe that the situation is rather complicated! We note that  $0 \in X$  and  $f(0) = 0$ , furthermore, since  $X$  is a cone, we observe that if for any  $x \in X$  we had  $f(x) < 0$  then indeed  $f$  is unbounded below on  $X$ . Thus the only situation in which a minimum exists is when a minimizing  $x$  is  $x_0 = 0$ , and of course  $f$  does not have a well defined gradient at the origin except in case  $C = 0$ .

Pursuing the above line of reasoning let us assume that  $f(x) = ax + (x^T C x)^{\frac{1}{2}} \geq 0$  whenever  $x \in X$  and that, in addition, the minimum of  $f$  (i.e., zero) is taken on at some point  $x_0$  such that  $x_0^T C x_0 > 0$ , and thus the gradient of  $f$  is defined at  $x_0$ . Summarizing the preceding,  $x_0$  satisfies:

$$(2) \quad x_0 \in R^n, \quad Ax_0 \leq 0, \quad ax_0 + (x_0^T C x_0)^{\frac{1}{2}} = 0, \quad x_0^T C x_0 > 0.$$

Using (2) the condition that  $(x - x_0)^T f'(x_0) \geq 0$  whenever  $Ax \leq 0$  may be written as:

$$(3) \quad \left[ a + \frac{x_0^T C}{(x_0^T C x_0)^{\frac{1}{2}}} \right] x \geq 0 \quad \text{whenever } Ax \leq 0 .$$

Up to this point, we have assumed that  $x_0$  satisfies (2) and that  $0 = f(x_0) \leq f(x)$  whenever  $x \in X$ ; we wrote down (3) as a statement "to be contemplated" and have not said anything about the truth or falsity of (3). However, it is quite trivial to show that when  $x_0$  satisfies (2) and if (3) is true then, necessarily,  $f(x_0) \leq f(x)$  for all  $x \in X$ . That is, from the fact that

$$(4) \quad x_0^T C x \leq (x_0^T C x_0)^{\frac{1}{2}} (x^T C x)^{\frac{1}{2}}$$

which holds for any  $x_0, x \in R^n$  [5, Lemma 1] and (3) we conclude that  $x \in R^n$  and  $Ax \leq 0$  implies  $f(x) = ax + (x^T C x)^{\frac{1}{2}} \geq 0$ . The converse statement, that one can conclude (3) from (2) and the fact that  $f(x) \geq 0$  whenever  $x \in X$  follows from Theorem 2 and the fact that  $f$  is then differentiable and convex in some small neighborhood of  $x_0$ . We have thus obtained a characterization of a minimizing  $x_0$  which satisfies  $x_0^T C x_0 > 0$ . Before attempting to dispose of minimizing  $x_0$ 's satisfying  $x_0^T C x_0 = 0$  (and thus, by (4),  $Cx_0 = 0$ ) we observe that (3) is of the form:  $Ax \leq 0$  implies  $dx \leq 0$ , thus by Lemma 1 we know that (3) is equivalent to the statement:

$$(5) \quad \text{There exists } \pi \in R^m \text{ such that}$$

$$\pi \geq 0 \quad \text{and} \quad \pi^T A + a + \frac{x_0^T C}{(x_0^T C x_0)^{\frac{1}{2}}} = 0 .$$



We observe next that if there exists an  $x_0$  satisfying (2) and (5) then by a suitable normalization we should be able to assume  $x_0^T C x_0 = 1$ ; specifically, letting  $z = (x_0^T C x_0)^{-\frac{1}{2}} x_0$  we obtain from (2) and (5) the conditions:

$$(6) \quad \begin{aligned} z &\in R^n, \quad \pi \in R^m \\ Az &\leq 0, \quad \pi \geq 0 \\ \pi^T A + a + z^T C &= 0, \quad z^T C z = 1. \end{aligned}$$

Again, by using (4), it is trivial to show that if there exist  $\pi$  and  $z$  satisfying (6) then  $f(x) \geq 0$  whenever  $x \in X$ . In fact, and this is very crucial to our development, we note that the same is true if we relax (6) to read:

$$(6') \quad \begin{aligned} z &\in R^n, \quad \pi \in R^m \\ Az &\leq 0, \quad \pi \geq 0 \\ \pi^T A + a + z^T C &= 0, \quad z^T C z \leq 1. \end{aligned}$$

We note that (6') is a more "realistic" statement in view of the fact that the  $z$  needed to satisfy (6) may yield  $Cz = 0$ . That is, we should really like to conclude (6') from the fact that  $ax + (x^T C x)^{\frac{1}{2}} \geq 0$  whenever  $x \in R^n$  and  $Ax \leq 0$ ; this is indeed the case and follows directly from:

### THEOREM 3

Assumption: (i)  $C$ ,<sup>†</sup> is a real symmetric  $n \times n$  matrix,  $A$  is an  $m \times n$  real matrix,  $u^T \in R^n$ .  
(ii)  $(ux)^2 \leq x^T \Gamma x$  whenever  $x \in R^n$  and  $ux \geq 0$ ,  $Ax \leq 0$ .  
Conclusion: There exist  $z \in R^n$ ,  $\pi \in R^m$  such that  $\pi \geq 0$ ,  $Az \leq 0$ ,  
 $u = \pi^T A + z^T C$ ,  $z^T C z \leq 1$ .

The proof of Theorem 3 will be found in Appendix B.

<sup>†</sup>Not necessarily positive semi-definite.

0426485

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
																																0426485																											

### GENERAL PURPOSE SUMMARY CARD

AF FORM 93A 1 AUG 54

[illegible]

As mentioned above, a direct consequence of Theorem 3 is:

Lemma 2: Let  $f$  and  $X$  be as in (1), then  $f(x) \geq 0$  for all  $x \in X$  if, and only if, there exist  $\pi$  and  $z$  satisfying (6').

Proof: The sufficiency of (6') is, as outlined above, quite clear. Assuming that for each  $x \in X$  we have  $f(x) \geq 0$  and letting  $u = -a$ , we see that the assumptions of Theorem 3 hold while the conclusion of Theorem 3 is precisely (6').

### III. A Nonhomogeneous Problem

Let us modify (1) so that  $X$  is defined by the inequalities  $Ax \leq b$ , rather than by  $Ax \leq 0$ , where  $b$  is a fixed vector in  $R^m$ . First of all, it need no longer be true that  $x_0 = 0 \in X$ , in fact  $X$  may be empty and to dispose of this difficulty we shall assume the contrary, that there exists on  $x \in R^n$  such that  $Ax \leq b$  or equivalently (according to Lemma 1) that:

$$(7) \quad \text{There is no } \pi \in R^m \text{ such that } \pi \geq 0, \pi^T A = 0, \pi^T b < 0.$$

Since  $X$  need no longer be a cone, the minimum of  $f(x) = ax + (x^T C x)^{\frac{1}{2}}$  on  $X$  (when it exists) may be any real number; let us assume that  $M$  is a lower bound of  $f$  on  $X$ , i.e.,

$$(8) \quad ax + (x^T C x)^{\frac{1}{2}} \geq M \quad \text{whenever } x \in R^n \text{ and } Ax \leq b.$$

With  $b = 0$  and  $M = 0$  we would be in the situation discussed in Section II with Lemma 2 applicable. We shall see now that the more general condition (8) may also be characterized in a fashion analogous to the homogeneous case. Towards this let us consider the relations:

$$\begin{aligned}
(9) \quad & z \in R^n, \quad \pi \in R^m, \quad \lambda \in R \\
& \pi \geq 0, \quad Az \leq \lambda b \\
& \pi^T A + a + z^T C = 0, \quad \pi^T b + M \leq 0, \quad z^T C z \leq 1,
\end{aligned}$$

and:

#### THEOREM 4

The statement (8) is true if, and only if, there exist  $z$ ,  $\pi$  and  $\lambda$  satisfying (9).

Proof: If  $\pi$ ,  $z$ ,  $\lambda$  satisfy (9) and  $x \in R^n$  is such that  $Ax \leq b$  then:

$$\begin{aligned}
0 &= \pi^T Ax + ax + z^T Cx \leq \pi^T b + ax + z^T Cx \\
&\leq -M + ax + (z^T C z)^{\frac{1}{2}} (x^T C x)^{\frac{1}{2}} \\
&\leq -M + ax + (x^T C x)^{\frac{1}{2}}.
\end{aligned}$$

and thus  $M \leq ax + (x^T C x)^{\frac{1}{2}}$ .

Conversely, let us assume that (8) is true. We assert that the statement

$$\begin{aligned}
(10) \quad & \text{If } x \in R^n, \quad \eta \in R \text{ are such that} \\
& Ax - \eta b \leq 0, \quad \eta \geq 0 \text{ then } \eta M \leq ax + (x^T C x)^{\frac{1}{2}},
\end{aligned}$$

is true. Clearly, when  $\eta > 0$  (10) holds, simply consider  $\eta^{-1}x$  and compare (10) with (8) which is assumed true. In case  $\eta = 0$ , we should like to conclude  $ax + (x^T C x)^{\frac{1}{2}} \geq 0$  from  $Ax \leq 0$ ; the last follows from the facts that (i) the statement (7) is true and (ii)  $ax + (x^T C x)^{\frac{1}{2}}$  is convex, homogeneous and bounded below on the set  $\{x | Ax \leq b\}$ . We next note that (10) is precisely analogous to the homogeneous statement in Lemma 2, applying Lemma 2 we see that from the fact that (10) holds it follows that there must exist  $\pi$ ,  $z$  and  $\lambda$  satisfying (9).

Q.E.D.

#### IV. Concluding Remarks

The following remarks are in order:

- A. Lemma 2 is precisely the principal theorem of [5] where it is demonstrated directly by using a certain "pseudo-norm" on the set of sequences of points in  $R^n$ , together with Lemma 1. Here Lemma 2 depends ultimately on the theorem of Frank-Wolfe [7] that any quadratic function bounded below on a polygonal convex set achieves its minimum.
- B. A significant interpretation of Lemma 2 is to think of it as a direct generalization of the Minkowski-Farkas theorem (our Lemma 1), because the former reduces to the latter when  $C = 0$ .
- C. One can think of Lemma 2 as expressing the fact that a certain convex set is closed. The set  $T$  in question consists of all negatives of  $a$ 's for which there exist  $z$  and  $\pi$  satisfying (6'), these in turn are the tangent planes of the convex function  $g(x) = (x^T C x)^{\frac{1}{2}}$  over the cone  $\{x | Ax \leq 0\}$ . In general, for arbitrary convex (and even continuous)  $g$  this set of tangents need not be closed; e.g., if  $g$  exhibits "asymptotic" behavior, a limit of tangents need not be a tangent.
- D. In view of the preceding and also because of the relation of Lemma 2 to Theorem 2, it would be of interest to generalize Lemma 2 to a larger class of pairs  $(X, f)$ , e.g., keeping  $X$  a finite cone and allowing  $f$  to take the form  $f(x) = ax + \sum_{k=1}^p (x^T C_k x)^{\frac{1}{2}}$ , where each  $C_k$  is positive semi-definite.

# APPENDIX A

## Proof of Theorem 1

All quantities are as defined in Theorem 1. We first demonstrate the simpler half of the proof that if

$$(11) \quad f(x) - f(y) \geq (x - y)^T f'(y) \quad \text{whenever } x, y \in X,$$

then  $f$  is convex. Suppose  $u, v$  are elements of  $X$  and  $\lambda \in [0, 1]$ , let  $x = u$  and  $y = \lambda u + (1 - \lambda)v$  then from (11) we get:

$$(12) \quad f(\lambda u + (1 - \lambda)v) \leq f(u) - (1 - \lambda)(u - v)^T f'(y).$$

Similarly, letting  $x = v$  and  $y = \lambda u + (1 - \lambda)v$  we obtain from (11):

$$(13) \quad f(\lambda u + (1 - \lambda)v) \leq f(v) + \lambda(u - v)^T f'(y).$$

Multiplying (12) by  $\lambda$ , and (13) by  $(1 - \lambda)$ , then adding, we obtain the required inequality.

We show next that if  $f$  is convex on  $X$  then (11) must hold. First, we show the above true for  $n = 1$ , then reduce the general case to the one dimensional one. Let us assume that  $K$  is an open subset of  $\mathbb{R}$  and that  $f$  is differentiable on  $K$ , then the convexity of  $f$  on a convex subset  $X$  of  $K$  is simply the statement that:

$$(14) \quad f(\beta) \leq \left( \frac{\gamma - \beta}{\gamma - \alpha} \right) f(\alpha) + \left( \frac{\beta - \alpha}{\gamma - \alpha} \right) f(\gamma)$$

whenever  $\alpha < \beta < \gamma$  and  $\alpha, \gamma \in X$ .

It is readily seen that (14) is equivalent to:

$$(15) \quad \frac{f(\beta) - f(\alpha)}{\beta - \alpha} \leq \frac{f(\gamma) - f(\beta)}{\gamma - \beta}$$

whenever  $\alpha < \beta < \gamma$  and  $\alpha, \gamma \in X$ .

Letting first  $\beta$  approach  $\gamma$  and then, independently, letting  $\beta$  approach  $\alpha$  we obtain from (15):

$$(16) \quad \left. \begin{aligned} \frac{f(\gamma) - f(\alpha)}{\gamma - \alpha} &\leq f'(\gamma) \\ f'(\alpha) &\leq \frac{f(\gamma) - f(\alpha)}{\gamma - \alpha} \end{aligned} \right\} \quad \text{if } \alpha < \gamma \text{ and } \alpha, \gamma \in X.$$

which is in fact equivalent to (15). Lastly, we note that (16) is equivalent to (11). We just demonstrated that if  $n = 1$  and  $f$  satisfies the assumptions of Theorem 1 then (11) must hold whenever  $f$  is convex.

Finally, suppose  $n$  is any positive integer and  $f$  is convex on  $X$  and satisfies the assumptions of Theorem 1. Since  $K$  is open, for fixed  $x, y \in X \subset K$ , there must exist an open interval  $I_{xy}$  containing  $[0, 1]$  and such that  $\lambda x + (1 - \lambda)y$  is in  $K$  whenever  $\lambda$  is in  $I_{xy}$ . Defining the function  $h_{xy}$  on  $I_{xy}$  by

$$h_{xy}(\lambda) = f(\lambda x + (1 - \lambda)y) \quad , \quad \text{for } \lambda \in I_{xy} \quad ,$$

one checks immediately that  $h_{xy}$  must be convex on  $[0, 1]$ . Consequently, since Theorem 1 holds for  $n = 1$ , we know that:

$$(17) \quad h_{xy}(1) - h_{xy}(0) \geq (1 - 0)h'_{xy}(0)$$

which is precisely the statement (11).

Q.E.D.

It should be noted that in the first half of the above proof no use was made of the properties of the gradient of  $f$ , thus, in fact, the following theorem is true.

#### THEOREM 5

Let  $f$ ,  $K$  and  $X$  satisfy the assumption of Theorem 1, then  $f$  is convex on  $X$  if, and only if, there exists a function  $G : K \rightarrow R^n$  and such that

$$f(x) - f(y) \geq (x - y)^T G(y) \quad \text{for all } x, y \in X.$$

Similarly, one notes that the following is true:

#### THEOREM 6

Let  $K$  be a nonempty open convex subset of  $R^n$ ,  $f : K \rightarrow R$ , then  $f$  is convex and differentiable on  $K$  if, and only if, there exists a continuous function  $G : K \rightarrow R^n$  such that

$$f(x) - f(y) \geq (x - y)^T G(y) \quad \text{for all } x, y \in K.$$



APPENDIX B  
Proof of Theorem 3

Consider the linear inequalities:

$$(18) \quad \begin{aligned} z &\in R^n, \quad \pi \in R^m \\ \pi &\geq 0, \quad Az \leq 0, \quad u^T = A^T \pi + Cz, \end{aligned}$$

and let  $P$  consist of all ordered pairs  $(\pi, z)$  satisfying (18). We show first that  $P$  is nonempty. Since (18) represents a system of linear inequalities, the fact that  $P$  is empty is equivalent, by Lemma 1, to the fact that there exist  $x$  and  $y$  satisfying:

$$(19) \quad \begin{aligned} x &\in R^n, \quad y \in R^m \\ y &\geq 0, \quad Ax \leq 0, \quad Cx = A^T y, \quad ux > 0. \end{aligned}$$

Thus, if  $P$  were empty, we would get from (19):

$$x^T Cx = \pi^T Ax \leq 0,$$

consequently  $x^T Cx \leq 0$  which together with  $ux > 0$  and  $Ax \leq 0$  contradicts the assumption of Theorem 3. Thus  $P$  is nonempty.

Next, if for some  $(\pi, z) \in P$  we had  $z^T Cx \leq 1$  then we would have the desired conclusion of Theorem 3; assume that  $z^T Cz > 1$  whenever  $(\pi, z) \in P$ . Applying the result in [7] that a quadratic function defined on a polyhedral convex set attains its minimum, we know that there exist  $(\pi_0, z_0) \in P$  such that  $1 < z_0^T Cz_0 \leq z^T Cz$  for each  $(\pi, z) \in P$ . Furthermore, it is clear that

$$(20) \quad (z_0^T C)z_0 \leq (z_0^T C)z \quad \text{whenever } (\pi, z) \in P$$

Applying Lemma 1 to (20) (essentially, making use of duality in linear programming), we see that there must then exist  $\pi$  and  $x$  satisfying:

$$\begin{aligned}
 (21) \quad & x \in R^n, \quad \pi \in R^m \\
 & Ax \leq 0, \quad \pi \geq 0, \quad Az_0 \leq 0 \\
 & u^T = A^T \pi_0 + Cz_0, \quad Cx = A^T \pi + Cz_0, \quad ux = z_0^T Cz_0.
 \end{aligned}$$

The following relations then are consequences of (21):

$$\begin{aligned}
 z_0^T Cz_0 &= z_0^T Cx - z_0^T A^T \pi \\
 &= x^T Cz_0 - \pi^T Az_0 \\
 &= ux - \pi_0^T Ax - \pi^T Az_0 \\
 &\geq ux \\
 &= z_0^T Cz_0.
 \end{aligned}$$

As a result,  $\pi_0^T Ax = \pi^T Az_0 = 0$  and  $z_0^T Cz_0 = ux = x^T Cz_0$ . However,  $Ax \leq 0$  and  $ux > 1$ ; thus by assumption in Theorem 3 we have  $0 < (ux)^2 \leq x^T Cx$  and:

$$\begin{aligned}
 (x^T Cx)^{\frac{1}{2}} &\geq ux \\
 &= z_0^T Cz_0 \\
 &= x^T Cz_0 \\
 &= x^T Cx - \pi^T Ax \\
 &\geq x^T Cx.
 \end{aligned}$$

From the last relations we get  $x^T Cx \leq 1$  and thus  $ux \leq (x^T Cx)^{\frac{1}{2}} \leq 1$ , a contradiction.

Q. E. D.

## BIBLIOGRAPHY

- [ 1 ] R. Bellman, Introduction to Matrix Analysis, McGraw-Hill, New York, 1960.
- [ 2 ] A. Charnes and W. W. Cooper, "The Strong Minkowski-Farkas-Weyl Theorem for Vector Spaces over Ordered Fields," Proc. Natl. Acad. Sci., Vol. 44, (1958) p. 914-916.
- [ 3 ] G. B. Dantzig, Linear Programming and Extensions, Princeton University Press, Princeton, N. J. (In press).
- [ 4 ] E. Eisenberg, "A Note on Semi-Definite Matrices," Research Report No. 9, Operations Research Center, University of California, Berkeley, July 20, 1961.
- [ 5 ] E. Eisenberg, "Supports of a Convex Function," Bull. Am. Math. Soc., Vol. 68, No. 3, (May 1962) p. 192-195.
- [ 6 ] Ky Fan, "Minimax Theorems," Proc. Natl. Acad. Sci., Vol. 39, No. 1, (January 1953) p. 42-47.
- [ 7 ] M. Frank and P. Wolfe, "An Algorithm for Quadratic Programming," Nav. Res. Log. Quart., Vol. 3, (March and June 1956) p. 95-110.
- [ 8 ] S. Karlin, Mathematical Methods and Theory in Games, Programming and Economics, Addison-Wesley Press, Cambridge, Massachusetts, 1959.
- [ 9 ] H. Markowitz, Portfolio Selection - An Efficient Diversification of Investments, Wiley Press, New York, 1959.

BASIC DISTRIBUTION LIST  
FOR UNCLASSIFIED TECHNICAL REPORTS

Head, Logistics and Mathematical  
Statistics Branch  
Office of Naval Research  
Washington 25, D. C.

C. O., ONR Office  
Navy No. 100, Box 39, F. P. O.  
New York, New York

ASTIA Document Service Center  
Building No. 5  
Cameron Station  
Alexandra, Virginia

Institute for Defense Analyses  
Communications Research Division  
von Neumann Hall  
Princeton, New Jersey

Technical Information Officer  
Naval Research Laboratory  
Washington 25, D. C.

C. O., ONR Branch Office  
1030 East Green Street  
Pasadena 1, California  
ATTN: Dr. A. R. Laufer

Bureau of Supplies and Accounts  
Code OW, Department of the Navy  
Washington 25, D. C.

Professor Russell Ackoff  
Operations Research Group  
Case Institute of Technology  
Cleveland 6, Ohio

Professor Kenneth J. Arrow  
Serra House, Stanford University  
Stanford, California

Professor G. L. Bach  
Carnegie Institute of Technology  
Planning and Control of Industrial  
Operations, Schenley Park  
Pittsburgh, 13, Pennsylvania

Professor L. W. Cohen  
Mathematics Department  
University of Maryland  
College Park, Maryland

Professor Donald Eckman  
Director, Systems Research Center  
Case Institute of Technology  
Cleveland, Ohio

Professor Lawrence E. Fouraker  
Graduate School of Business  
Harvard University  
Cambridge, Massachusetts

Professor David Gale  
Department of Mathematics  
Brown University  
Providence 12, Rhode Island

Dr. Murray Geisler  
The RAND Corporation  
1700 Main Street  
Santa Monica, California

Professor L. Hurwicz  
School of Business Administration  
University of Minnesota  
Minneapolis 14, Minnesota

Dr. James R. Jackson, Director  
Western Management Sciences Institute  
University of California  
Los Angeles 24, California

Professor Samuel Karlin  
Mathematics Department  
Stanford University  
Stanford, California

Professor C. E. Lemke  
Department of Mathematics  
Rensselaer Polytechnic Institute  
Troy, New York

Professor W. H. Marlow  
Logistics Research Project  
The George Washington University  
707 - 22nd Street, N. W.  
Washington 7, C. D.

BASIC DISTRIBUTION LIST  
FOR UNCLASSIFIED TECHNICAL REPORTS

Professor A. Charnes  
The Technological Institute  
Northwestern University  
Evanston, Illinois

Professor R. Radner  
Department of Economics  
University of California  
Berkeley 4, California

Professor Murray Rosenblatt  
Department of Mathematics  
Brown University  
Providence 12, Rhode Island

Mr. J. R. Simpson  
Bureau of Supplies and Accounts  
Navy Department (Code W31)  
Washington 25, D. C.

Professor A. W. Tucker  
Department of Mathematics  
Princeton University  
Princeton, New Jersey

Professor J. Wolfowitz  
Department of Mathematics  
Lincoln Hall, Cornell University  
Ithaca 1, New York

C. O., ONR Branch Office  
346 Broadway  
New York 13, New York  
ATTN: J. Laderman

Professor Oskar Morgenstern  
Economics Research Project  
Princeton University  
92 A Nassau Street  
Princeton, New Jersey